

P-ADIC FUNCTORIALITY FOR INNER FORMS OF UNITARY GROUPS IN THREE VARIABLES

JUDITH LUDWIG

ABSTRACT. We prove p -adic functoriality for inner forms of unitary groups in three variables by establishing the existence of morphisms between eigenvarieties that extend the classical Langlands functoriality.

1. INTRODUCTION

Eigenvarieties are rigid analytic spaces that p -adically interpolate systems of Hecke eigenvalues attached to automorphic representations. For definite unitary groups they have been constructed by Chenevier [Che04] using Buzzard’s and Coleman’s approach [Buz07] and by Emerton [Eme06] using completed cohomology. Given a construction of eigenvarieties for a class of groups a question that naturally arises is whether one can extend Langlands functoriality p -adically. The aim is to construct a morphism of eigenvarieties that on classical points agrees with the classical transfer. Chenevier was able to prove p -adic functoriality in the case of the Jacquet–Langlands transfer by interpolating the classical transfer [Che05]. Using his approach White [Whi12] established a p -adic endoscopic transfer for unitary groups. Flicker [Fli10] proved a p -adic interpolation of base-change. Newton [New12] proved a p -adic Jacquet–Langlands correspondence in Emerton’s setting.

In these notes we prove p -adic functoriality for certain inner forms G and G' of a quasi-split unitary group U_3^* in three variables. There are various problems one encounters when trying to interpolate the classical transfer:

- The classical transfer can only be proved for L -packets (or A -packets) rather than for single automorphic representations and in general the multiplicities of elements in a packet may vary and can even be zero yet eigenvarieties parameterize systems of eigenvalues of Hecke operators that show up in certain spaces of p -adic automorphic forms.
- An eigenvariety comes with a tame level. If we want to construct a p -adic functoriality morphism from an eigenvariety for G that has arbitrary tame level to an eigenvariety for G' , we need to find a suitable tame level for G' .

We deal with these problems as follows:

We study “stable” situations, i.e. situations where the multiplicities within an L -packet are constant. This allows us to transfer automorphic representations rather than packets. In order to find suitable tame levels of G and G' that are respected by the classical transfer, we use the theory of the Bernstein decomposition and properties of the local Jacquet–Langlands transfer.

Our main theorem is Theorem 4.2, where we prove the existence and uniqueness of the p -adic functoriality morphism. We remark that the source can be an eigenvariety for G that has *arbitrary* tame level at the bad primes, i.e. the primes, where

G and G' differ. In this aspect these notes differ from the papers on p -adic functoriality cited above, where the open compact subgroup at such primes is usually assumed to be maximal.

Acknowledgements. I would like to thank Kevin Buzzard for suggesting that I think about this topic. I would like to thank him and Paul-James White for many helpful conversations and their comments on an earlier draft of this paper. I am very grateful to Sug Woo Shin for helpful correspondence and in particular for explaining the arguments needed for the multiplicity statement in section 2. Moreover I thank Christian Johansson for many helpful conversations. I would like to thank Imperial College for providing a Roth Studentship and EPSRC for their financial support of my doctoral studies.

Notation. All division algebras occurring in this paper will be central simple algebras.

2. CLASSICAL FUNCTORIALITY

Let E/\mathbb{Q} be an imaginary quadratic extension and U_3^* the quasi-split unitary group in three variables attached to E/\mathbb{Q} . Let G, G' be two definite unitary groups that are inner forms of U_3^* . Note that for all places p that are non-split in E the local groups G_p and G'_p are quasi-split and therefore isomorphic. Assume that G comes from a division algebra D . We denote by S the set of places p where $G_p \not\cong G'_p$ and assume furthermore that $G'_p \cong \mathrm{GL}_3/\mathbb{Q}_p$ for all places $p \in S$. Denote by JL the local Jacquet–Langlands transfer (see Théorème principal in [DKV84], p.34).

We define S_G to be the set of all primes p that split in E and such that D_w is ramified for $w|p$ and similarly for G' . We fix inner automorphisms $\psi_G : G_{\overline{\mathbb{Q}}} \rightarrow (U_3^*)_{\overline{\mathbb{Q}}}$ and $\psi_{G'} : G'_{\overline{\mathbb{Q}}} \rightarrow (U_3^*)_{\overline{\mathbb{Q}}}$. Let $\Pi(G)$ (resp. $\Pi(G')$, $\Pi(U_3^*)$) denote the set of global L -packets of G (resp. G' , U_3^*) as defined in [Rog90] section 14.6. (cf. sections 12.2 and 14.4 for the definition of local L -packets). We would like to stress that for the transfer to work Rogawski has to enlarge certain local L -packets to what he calls A -packets (see [Rog90] p.199) and by definition the representations in a global L -packets vary locally within a local L -packet or an A -packet.

In section 14.4 and 14.6 Rogawski constructs transfer maps $\psi_G : \Pi(G) \rightarrow \Pi(U_3^*)$ and $\psi_{G'} : \Pi(G') \rightarrow \Pi(U_3^*)$. As G comes from a division algebra all L -packets of G are stable, i.e. non-endoscopic (cf. [Rog90] Theorem 14.6.3 and p.201 for the definition of a stable L -packet), so using his notation we get $\Pi(G) = \Pi_s(G)$. Furthermore Rogawski proves the following proposition:

Proposition 2.1. *The map ψ_G defines a bijection between $\Pi_s(G)$ and*

$$Im(\psi_G) := \{\Pi \in \Pi_s(U_3^*) : \dim(\Pi) = 1 \text{ or } \Pi_v \in \Pi^2(U_{3,v}^*) \text{ for all } v \in S_G \cup \{\infty\}\}.$$

Proof. This is Proposition 14.6.2 in [Rog90]. \square

Lemma 2.2. *Let H be an inner form of U_3^* . Assume $\Pi \in \Pi_s(H)$ is a stable L -packet. Then*

$$m(\pi) = 1 \text{ for all } \pi \in \Pi.$$

Proof. By [Rog90] equation (14.6.2) and the proof of Proposition 14.6.2 we get the following identity of distributions

$$\sum_{\pi \in \Pi} m(\pi) \mathrm{tr}(\pi(f)) = \mathrm{tr}(\psi_H(\Pi)(f^*)),$$

where f^* denotes the transfer of the function f (see [Rog90] section 14.2). Using the fact that the transfer ψ_H satisfies the character identity $\text{tr}(\Pi(f)) = \text{tr}(\psi_H(\Pi)(f^*))$ (see section 14.4) we get

$$\sum_{\pi \in \Pi} m(\pi) \text{tr}(\pi(f)) = \text{tr}(\Pi(f)) = \sum_{\pi \in \Pi} \langle 1, \pi \rangle \text{tr}(\pi(f)).$$

But $\langle 1, \pi \rangle \in \{\pm 1\}$ and $m(\pi) \geq 0$, so we deduce that

$$m(\pi) = 1 \quad \text{for all } \pi \in \Pi.$$

□

Theorem 2.3. *Let $\pi = \otimes \pi_p$ be an automorphic representation of G which is not one-dimensional. Then*

$$\pi' \cong \bigotimes_{p \in S} JL(\pi_p) \otimes \bigotimes_{p \notin S} \pi_p$$

is an automorphic representation of G' occurring with the same multiplicity as π .

Proof. Our assumptions on G and G' imply that $S_{G'} \subset S_G$ and so $\text{Im}(\psi_G) \subset \text{Im}(\psi_{G'})$. Therefore π' is an element of a stable L -packet. We can therefore apply the previous lemma to conclude that π' is automorphic. □

Following [Rog90] as in section 14.6 we can also transfer one-dimensional representations: Let $\mathbb{A}_{E,1}$ denote the subgroup of norm one elements in \mathbb{A}_E^* , the ideles in E , and E_1 the norm one elements in E^* . Let $N_G : G(\mathbb{A}) \rightarrow \mathbb{A}_{E,1}$ be defined by $N_G(x) = \prod_v N_{G,v}(x_v)$ where $N_{G,v}$ is the determinant for $v \notin S_G$ and the reduced norm Nrd for $v \in S_G$. If π is a one dimensional automorphic representation of G , then $\pi = \chi \circ N_G$, where χ is a character of $\mathbb{A}_{E,1}$ trivial on E_1 . The transfer of π is the one-dimensional representation $\pi' = \chi \circ N_{G'}$ of G' associated to this character.

Next we explore some properties of the local Jacquet–Langlands transfer that are needed to establish p -adic functoriality. We make use of the Bernstein decomposition, the notation is as in [BK98]: Let F/\mathbb{Q}_p be a finite extension with ring of integers \mathcal{O}_F and uniformizer ϖ , A/F a central simple algebra and A^* the group of units, so $A^* \cong \text{GL}_m(D)$ for D/F some division algebra. Let $d \in \mathbb{N}$ be such that $d^2 = \dim D$. The reduced Norm map will be denoted by $\text{Nrd} : A^* \rightarrow F^*$. Let O_D be a maximal order and ϖ_D a uniformizer in O_D . The subgroups

$$K_n := \{x \in \text{GL}_m(O_D) : x \equiv 1 \pmod{\varpi_D^n}\}$$

form a fundamental system of compact open neighbourhoods of the identity. Let $\mathcal{B}(A^*)$ be the set of Bernstein components of A^* . For $s \in \mathcal{B}(A^*)$ let $\mathcal{R}^s(A^*)$ be the category of isomorphism classes of smooth representations of A^* associated to s , which we will also call a Bernstein component. Let $\text{Irr } \mathcal{R}^s(A^*) \subset \mathcal{R}^s(A^*)$ be the class of objects that are irreducible representations.

For K a compact open subgroup of A^* we write

$$e_K := \mu(K)^{-1} \mathbf{1}_K \in C_c^\infty(A^*, \mathbb{C}),$$

for the idempotent attached to K , here $\mathbf{1}_K$ is the characteristic function on K and μ a Haar measure on A^* .

Now let $e \in C_c^\infty(A^*, \mathbb{C})$ be an arbitrary idempotent. Then there exists a compact open subgroup $K \subset A^*$ such that $e_K \cdot e = e = e \cdot e_K$, where “ \cdot ” denotes convolution. We choose K to be of the form K_n .

We denote by $\text{Irr } \mathcal{R}_e(A^*)$ the class of all irreducible smooth representations π such that $e \cdot \pi \neq 0$, i.e. of those irreducible representations that are generated by $e \cdot \pi$. Note that for any such representation π the set of K -fixed vectors is nonzero as

$$0 \neq e \cdot \pi = e \cdot e_K \cdot \pi = e \cdot \pi^K.$$

It follows from [Ber84] 3.9 that there exists a finite set $\mathcal{J}_e \subset \mathcal{B}(A^*)$ such that

$$\text{Irr } \mathcal{R}_e(A^*) \subset \bigcup_{s \in \mathcal{J}_e} \text{Irr } \mathcal{R}^s(A^*).$$

Proposition 2.4. *Let $e \in C_c^\infty(A^*, \mathbb{C})$ be an idempotent with the property that we can find a set \mathcal{J}_e as above such that any $s \in \mathcal{J}_e$ is associated to a supercuspidal representation. Then there exists a compact open subgroup K_e of $\text{GL}_{md}(F)$ such that*

$$JL(\pi)^{K_e} \neq 0$$

for all $\pi \in \text{Irr } \mathcal{R}_e(A^*)$.

Proof. Let $\pi \in \text{Irr } \mathcal{R}_e(A^*)$. There exists $s_\pi \in \mathcal{J}_e$ such that $\pi \in \text{Irr } \mathcal{R}^{s_\pi}(A^*)$. For $s \in \mathcal{J}_e$ let $\rho_s \in \text{Irr } \mathcal{R}^s(A^*)$ be a representative. Then π is an unramified twist of ρ_{s_π} , say $\pi \cong \rho_{s_\pi} \otimes \chi$. Note that χ is trivial on K and therefore $\dim(\pi^K) = \dim(\rho_{s_\pi}^K) \neq 0$. So for a general $\pi \in \text{Irr } \mathcal{R}_e(A^*)$ we have

$$\dim(\pi^K) \in \{\dim(\rho_s^K) : s \in \mathcal{J}_e\}.$$

The fact that χ is unramified implies that for any $K' \subset \text{GL}_{md}(\mathcal{O}_F)$ open compact

$$\dim(JL(\pi)^{K'}) \in \{\dim(JL(\rho_{s_\pi})^{K'}) : s \in \mathcal{J}_e\},$$

as the Jacquet–Langlands transfer respects twists by a character. It is therefore sufficient to find an open compact open subgroup $K_e \subset \text{GL}_{md}(\mathcal{O}_F)$ such that $JL(\rho_{s_\pi})^{K_e} \neq 0$ for all $s \in \mathcal{J}_e$. Such a K_e exists as $JL(\rho) = \bigcup_K JL(\rho)^K$ and \mathcal{J}_e is finite. \square

Remark 2.5. *Assume that $m = 1$, so that $A^* = D^*$. As D^* has no proper Levi subgroups defined over F all irreducible smooth representations are supercuspidal, so we can apply Proposition 2.4 for any idempotent $e \in C_c^\infty(D^*, \mathbb{C})$.*

Lemma 2.6. *Let $\pi = \chi \circ \text{Nrd} : A^* \rightarrow F^*$ be a character of A^* invariant under K_n . Then $\pi' := \chi \circ \det : \text{GL}_{md}(F) \rightarrow F^*$ is invariant under $K'_n := \{g \in \text{GL}_{md}(\mathcal{O}_F) : \det(g) \in 1 + \varpi^n \mathcal{O}_F\}$.*

Proof. The lemma follows from the fact that $\text{Nrd}(K_n) = 1 + \varpi^n \mathcal{O}_F = \det(K'_n)$. \square

3. EIGENVARIETIES FOR DEFINITE UNITARY GROUPS

To set up notation we will briefly recall the definition and existence theorem of eigenvarieties of idempotent type as constructed in section 7 of [BC09]. Let E/\mathbb{Q} and G be as in section 2. Let p be a prime that splits in E and such that $G_p \cong \text{GL}_3/\mathbb{Q}_p$. Fix such an isomorphism. Furthermore fix embeddings $\iota_p : \overline{\mathbb{Q}} \rightarrow \overline{\mathbb{Q}}_p$ and $\iota_\infty : \overline{\mathbb{Q}} \rightarrow \mathbb{C}$.

Let S_1 be a finite set of places of \mathbb{Q} containing the archimedean place such that G is unramified outside S_1 . Fix a model G/\mathbb{Z}_{S_1} of G . Let $\mu = \prod_l \mu_l$ be a product Haar measure on $G(\mathbb{A}_f)$ such that $\mu_l(\mathbb{Z}_l) = 1$ for all $l \notin S_1$.

Let S_0 be a subset of the primes l that split in E and such that $G_l \cong \mathrm{GL}_n/\mathbb{Q}_l$. Assume furthermore that $p \notin S_0$ and $S_0 \cap S_1 = \emptyset$. Define

$$\mathcal{H} := \mathcal{A}_p \otimes \mathcal{H}_{\mathrm{ur}},$$

where $\mathcal{H}_{\mathrm{ur}} = \mathcal{C}(G(\mathbb{Z}_{S_0}) \backslash G(\mathbb{A}_{S_0})/G(\mathbb{Z}_{S_0}), \mathbb{Z})$ ¹ and \mathcal{A}_p is the commutative Atkin-Lehner subring of $C_c^\infty(I \backslash \mathrm{GL}_3(\mathbb{Q}_p)/I, \mathbb{Z}[\frac{1}{p}])$, where I denotes the standard Iwahori subgroup of $\mathrm{GL}_3(\mathbb{Q}_p)$ (see [BC09] section 6.4.1 for the definition). For $u = \mathrm{diag}(p^2, p, 1) \in \mathrm{GL}_3(\mathbb{Q}_p)$ let $[IuI] \in \mathcal{A}_p$ be the characteristic function on the double coset IuI . The element $[IuI] \otimes 1 \in \mathcal{H}$ will be denoted by u_0 .

Let $\mathcal{Z}_0 \subset \mathrm{Hom}_{\mathrm{ring}}(\mathcal{H}, \overline{\mathbb{Q}_p}) \times \mathbb{Z}^n$ be the set of pairs $(\psi_{(\pi, \mathcal{R})}, \underline{k})$ where (π, \mathcal{R}) runs over the set of p -refined automorphic representations, $\psi_{(\pi, \mathcal{R})}$ is the system of Hecke eigenvalues attached to (π, \mathcal{R}) and \underline{k} is the weight of π (see [BC09] section 7.2.2 for the definition of a p -refined automorphic representation and details regarding $\psi_{(\pi, \mathcal{R})}$). Let \mathcal{Z} be a subset of \mathcal{Z}_0 .

Weight space will be denoted by \mathcal{W} . It is the quasi-separated rigid analytic space over \mathbb{Q}_p

$$\mathcal{W} := \mathrm{Hom}_{\mathrm{cont}}((\mathbb{Z}_p^*)^n, \mathbb{G}_m),$$

whose points over any affinoid \mathbb{Q}_p -algebra A parameterize the continuous characters $(\mathbb{Z}_p^*)^n \rightarrow A^*$. We embed \mathbb{Z}^n into $\mathcal{W}(\mathbb{Q}_p)$ by

$$(k_1, \dots, k_n) \mapsto \left((t_1, \dots, t_n) \mapsto \prod_{i=1, \dots, n} t_i^{k_i} \right).$$

An eigenvariety for \mathcal{Z} is defined as follows (for a definition of an accumulation and Zariski-dense subset of a rigid space see [BC09] section 3.3.1).

Definition 3.1. *Let L be a finite extension of \mathbb{Q}_p . An eigenvariety (X, ψ, ν, Z) for \mathcal{Z} over L is a reduced p -adic rigid analytic space X over L together with*

- a ring homomorphism $\psi : \mathcal{H} \rightarrow \mathcal{O}(X)^{\mathrm{rig}}$
- an analytic map $\omega : X \rightarrow \mathcal{W}$ defined over L and
- an accumulation and Zariski-dense subset $Z \subset X(\overline{\mathbb{Q}_p})$

such that the following conditions are satisfied:

- (1) The map $\nu := (\omega, \psi(u_0)^{-1}) : X \rightarrow \mathcal{W} \times \mathbb{G}_m$ is finite.
- (2) For all open affinoid $V \subset \mathcal{W} \times \mathbb{G}_m$, the natural map

$$\psi \otimes \nu^* : \mathcal{H} \times \mathcal{O}(V) \rightarrow \mathcal{O}(\nu^{-1}(V))$$

is surjective.

- (3) The natural evaluation map $X(\overline{\mathbb{Q}_p}) \rightarrow \mathrm{Hom}_{\mathrm{ring}}(\mathcal{H}, \overline{\mathbb{Q}_p})$

$$x \mapsto \psi_x := (h \mapsto \psi(h)(x))$$

induces a bijection $Z \rightarrow \mathcal{Z}, z \mapsto (\psi_z, \omega(z))$.

Next we introduce the relevant sets \mathcal{Z} that give rise to the eigenvarieties we want to compare via functoriality. Fix an idempotent

$$e \in C_c^\infty(G(\mathbb{A}_f^{p, S_0}), \overline{\mathbb{Q}}) \otimes 1_{\mathcal{H}_{\mathrm{ur}}} \subset C_c^\infty(G(\mathbb{A}_f^p), \overline{\mathbb{Q}})$$

¹ If l is a prime which is inert in E , the extension E_v/\mathbb{Q}_l is unramified and as $G_l \times \mathrm{Spec}(E_v) \cong \mathrm{GL}_n$, the group G is unramified at l . In the definition of \mathcal{H} we could have included the (commutative) unramified Hecke algebras at these primes and all constructions could be made using this enlarged Hecke algebra.

and a finite extension L of \mathbb{Q}_p that contains the values of $i_p(e) := i_p \circ e$. Let $\mathcal{Z}_e \subset \mathcal{Z}_0$ be the subset of $(\psi_{(\pi, \mathcal{R})}, \underline{k})$ such that $e \cdot \pi^p \neq 0$.

Recall that a rigid space X over L is called *nested* if it has an admissible covering by open affinoids $\{X_i, i \geq 0\}$ such that $X_i \subset X_{i+1}$ and the natural L -linear map $\mathcal{O}(X_{i+1}) \rightarrow \mathcal{O}(X_i)$ is compact.

Theorem 3.2. *There exists a unique eigenvariety $(X/L, \psi, \nu, Z)$ for \mathcal{Z}_e . It is nested and equidimensional of dimension 3.*

Proof. This is Theorem 7.3.1 in [BC09]. The uniqueness assertion is proved in their Proposition 7.2.8.

4. p -ADIC TRANSFER OF AUTOMORPHIC FORMS

Let G and G' be unitary groups as in section 2. Fix models G/\mathbb{Z}_{S_1} and G'/\mathbb{Z}_{S_1} where we've chosen S_1 such that both groups are unramified outside S_1 . Just as above let S_0 be a subset of the primes l that split in E and such that $G_l \cong \mathrm{GL}_3/\mathbb{Q}_l$, so in particular $S_G \cap S_0 = \emptyset$, where S_G is as in section 2. Note that our assumptions on G' imply that for $l \in S_0$ we also have $G'_l \cong \mathrm{GL}_3/\mathbb{Q}_l$. Assume furthermore that $S_0 \cap S_1 = \emptyset$. Fix an idempotent

$$e \in C_c^\infty(G(\mathbb{A}_f^{p, S_0}), \overline{\mathbb{Q}}) \otimes 1_{\mathcal{H}_{\mathrm{ur}}} \subset C_c^\infty(G(\mathbb{A}_f^p), \overline{\mathbb{Q}})$$

and assume

$$(1) \quad e = \prod_{l \notin S_0 \cup \{p\}} e_l \times \prod_{l \in S_0} e_{G(\mathbb{Z}_l)}$$

is a product of local idempotents $e_l \in C_c^\infty(G(\mathbb{Q}_l), \overline{\mathbb{Q}})$.

As in section 2 we let S be the set of places where the local groups are not isomorphic. For $l \in S$ choose a compact open subgroup $K_l \subset G'(\mathbb{Q}_l)$ such that $(\pi'_l)^{K_l} \neq 0$ for all local components π'_l of automorphic representations π' of G' that are the transfer of an automorphic representation π of G satisfying $e \cdot \pi \neq 0$. The existence of such compact open subgroups is guaranteed by Proposition 2.4 and Lemma 2.6. Define the idempotent

$$e' := \prod_{l \in S} e_{K_l} \times \prod_{l \notin S \cup S_0 \cup \{p\}} e_l \times \prod_{l \in S_0} e_{G'(\mathbb{Z}_l)}.$$

Note that for any p -refined automorphic representation (π, \mathcal{R}) of G

$$\psi_{(\pi, \mathcal{R})} = \psi_{(\pi', \mathcal{R})},$$

where π' is the transfer of π . We have established an inclusion

$$(2) \quad \mathcal{Z}_e \subset \mathcal{Z}_{e'}.$$

Lemma 4.1. *Let $\mathcal{Z}_1 \subset \mathcal{Z}_2$ be two subsets of $\mathcal{Z}_0 \subset \mathrm{Hom}_{\mathrm{ring}}(\mathcal{H}, \overline{\mathbb{Q}}_p) \times \mathbb{Z}^n$. Assume there exist eigenvarieties $(X_1/L, \psi_1, \nu_1, Z_1)$ for \mathcal{Z}_1 and $(X_2/L, \psi_2, \nu_2, Z_2)$ for \mathcal{Z}_2 . Then there exists a unique closed immersion*

$$\zeta : X_1 \hookrightarrow X_2$$

over L such that $\nu_1 = \zeta \circ \nu_2$ and for all $h \in \mathcal{H}$, $\psi_1(h) = \zeta^*(\psi_2(h))$.

Proof. By Lemma 7.2.7 (b) of [BC09] and property 3 of an eigenvariety there exists a unique injection $\zeta : Z_1 \hookrightarrow Z_2$ such that for all $z \in Z_1$, $\psi_{1z} = \psi_{2_{\zeta(z)}}$ and $\omega_1(z) = \omega_2(\zeta(z))$. Our claim is that we can extend ζ to a closed immersion $\zeta : X_1 \hookrightarrow X_2$. Using the fact that the X_i are reduced and Lemma 7.2.7 (b) of [BC09] again

we see that such an extension if it exists is unique. The proof of the existence is contained verbatim in the proof of the uniqueness of eigenvarieties, that is in the proof of proposition 7.2.8 of [BC09]. More precisely they construct a closed immersion $X_1 \hookrightarrow X_2$. The same construction applies in our situation and proves the Lemma. \square

Theorem 4.2. *Let G, G', S_0, e and e' be as above. Let L/\mathbb{Q}_p be a finite extension that contains the values of $i_p(e)$ and of $i_p(e')$.*

Let $(X/L, \psi, \nu, Z_e)$ (resp. $(X'/L, \psi', \nu', Z'_e)$) be the eigenvariety for Z_e (resp. $Z_{e'}$). Then there exists a unique closed immersion of eigenvarieties

$$\zeta : X \hookrightarrow X'$$

compatible with the inclusion 2, i.e. with the classical transfer of automorphic representations. The image of ζ is a union of irreducible components.

Proof. Everything except the last assertion follows from the previous lemma. The last assertion follows from [Con99] Corollary 2.2.7, which states that for a rigid space X , which is equidimensional of dimension m , the analytic subsets in X , which are equidimensional of dimension m , are exactly the union of irreducible components. \square

REFERENCES

- [BC09] Joël Bellaïche and Gaëtan Chenevier. Families of Galois representations and Selmer groups. *Astérisque*, (324), 2009.
- [Ber84] J. N. Bernstein. Le “centre” de Bernstein. In *Representations of reductive groups over a local field*, Travaux en Cours, pages 1–32. Hermann, Paris, 1984.
- [BK98] Colin J. Bushnell and Philip C. Kutzko. Smooth representations of reductive p -adic groups: structure theory via types. *Proc. London Math. Soc.* (3), 77(3):582–634, 1998.
- [Buz07] Kevin Buzzard. Eigenvarieties. In *L-functions and Galois representations*, volume 320 of *London Math. Soc. Lecture Note Ser.*, pages 59–120. Cambridge Univ. Press, Cambridge, 2007.
- [Che04] Gaëtan Chenevier. Familles p -adiques de formes automorphes pour GL_n . *J. Reine Angew. Math.*, 570:143–217, 2004.
- [Che05] Gaëtan Chenevier. Une correspondance de Jacquet–Langlands p -adique. *Duke Math. J.*, 126(1):161–194, 2005.
- [Con99] Brian Conrad. Irreducible components of rigid spaces. *Ann. Inst. Fourier (Grenoble)*, 49(2):473–541, 1999.
- [DKV84] P. Deligne, D. Kazhdan and M.-F. Vignéras. Représentations des algèbres centrales simples p -adiques. In *Representations of reductive groups over a local field*, Travaux en Cours, pages 1–32. Hermann, Paris, 1984.
- [Eme06] Matthew Emerton. On the interpolation of systems of eigenvalues attached to automorphic Hecke eigenforms. *Invent. Math.*, 164(1):1–84, 2006.
- [Fli10] Yu. Z. Flicker. Base change for Hilbert eigenvarieties of unitary groups. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 377:148–198, 244, 2010.
- [New12] James Newton. Completed cohomology of Shimura curves and a p -adic Jacquet–Langlands correspondence. *Mathematische Annalen*, pages 1–35, 2012.
- [Rog90] Jonathan D. Rogawski. *Automorphic representations of unitary groups in three variables*, volume 123 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1990.
- [Whi12] Paul-James White. p -adic Langlands functoriality for the definite unitary group. *preprint*, arXiv:1206.2193v1, 2012.